


# Total Joint Replacement

**Hoag Orthopedic Institute  
Surgery Centers**

**Posterior Approach  
Total Hip Replacement**



A photograph of surfers in the ocean. In the foreground, a surfer in a black wetsuit is seen from behind, carrying a light blue surfboard and splashing water. In the background, two other surfers are visible, one with a red surfboard. The scene is set against a backdrop of mountains and a bright sky. The text "We Get You Back To You" is overlaid in the center in a white, sans-serif font.

We Get You Back To You

# The Hip

## Hip Joint

- Ball and socket joint made up of the:
  - Head of the femur (the “ball”)
  - Acetabulum (the “socket”)

## Ligament


- Tough, flexible fibrous connective tissue which connects two bones or cartilages or holds a joint together.

## Cartilage

- Strong, flexible connective tissue that protects your joints and bones.

## Synovial fluid

- Liquid that lubricates the joint and keeps it moving smoothly.



# Hip Replacement

The whole joint is removed and replaced with an artificial joint.

Components include:

- Acetabular component
- Plastic liner
- Femoral head
- Femoral stem



# Pre-Surgery: Preparation

## Review your surgery forms.

Make appointments for:

- Primary care physician for medical clearance
- Any ordered testing
  - Labs, EKG, etc.

Complete pre-operative forms.

## Complete the requirements outlined in your surgery packet.

- Return required information to your surgeon's office at least one week before your surgery, preferably two.



# Pre-Surgery: Plan Your Recovery

Start planning your recovery at home now.

Determine who will help you:

- Spouse, family member, friend

Determine where you will recovery:

- Your own home, at a family member or friend's home

Plan to have your caregiver be available to help for the first 48-72 hours at home.



# Pre-Surgery: Plan Your Recovery


## Clothing, the day of surgery and during your recovery.

- Wear loose, comfortable clothes that are easy to get on and off.
  - Shorts and dresses are often easier than long pants.
- Choose shoes with closed toes and backs that you can slip on and off.



# Pre-Surgery: Home Preparation

Safety  
Zone



Your “safety zone” is the area between your shoulders and hips. Keeping commonly used items in this area prevents you having to bend and reach thus reducing the risk of falls.

## Arrange

Arrange frequently used items in your “safety zone.”

## Remove

Remove loose throw rugs.

## Clear

Clear hallways for use of walker.

## Make

Make arrangements for pets.

## Use

Use night lights.

## Create

Create charging stations next to areas where you sit or lie down



# Pre-Surgery: Bathroom Preparation

Make sure you have non-skid surfaces.

## Purchase:

- 3:1 Commode (needed to maintain precautions)

## Equipment that may be helpful:

- Shower Chair
- Tub Transfer Bench
- Handheld shower nozzle



Compare prices online and at your local DME store.

# Pre-Surgery: Equipment

## Equipment to help you maintain your precautions:

- Reacher / Grabber
- Long Reach Sponge
- Long Reach Shoehorn
- Sock Aid



This equipment helps you maintain your posterior approach precautions.  
Consider purchasing equipment pre-operatively.

Compare prices online and at your local DME store.

# Pre-Surgery: Equipment

## Walkers

- Walkers are used after surgery (think safety)
- Your surgeon's office will tell you how to acquire a walker
- **If You are Purchasing a Walker:**
  - Amazon.com, Walmart.com, your local DME store
  - Your walker will come with instructions for adjusting the height. Please review the instructions for fit and adjustment before your surgery.
  - *If you are instructed to purchase a walker, contact your insurance company to see if you can be reimbursed.*

## A few things to remember about fit

- When your arms are relaxed at your sides:
  - The crease on the inside of your wrist should line up with the walker grips
- When holding onto the walker grips:
  - Relax your shoulders
  - Your elbows will be slightly bent



# Pre-Surgery: Practice Stairs

## Stairs:

- Patients can use stairs the day of surgery.
- If you have the option, plan to spend your first night or few nights downstairs.
- Practice before your surgery.



# Pre-Surgery: Practice Getting in and out of a Car

## Before getting into a car:

- Have someone move the seat as far back as it will go.
- Recline the back of the seat if possible.
- Remember your precautions.

## Getting into a car:

- Back up to the open doorway of the car.
- Hold onto the side of the car or dashboard and the walker for support.
- Lower yourself slowly onto the seat. Watch your head.
- Bringing your legs into the car by:
  - Sliding back into the center of the seat so you are not sitting on the edge.
  - Lifting your legs one at a time into the car.
- Keep your car seat as high as possible.

## Getting out of a car:

- To bring your legs out of the car.
  - Slide to the edge of the seat.
  - Lift your legs out of the car one at a time.
- To stand up
  - Make sure your feet are on the ground
  - Place the walker as close as possible
  - Use the car seat or jam (a car cane can help) and the crossbar of the walker to stand (never try to stand with just the walker, it could tip over)
  - Raise yourself slowly



# Pre-surgery: Optimization

## Nutrition: Before and After Surgery

Begin increasing protein intake a few weeks before surgery to build up strength and to help with healing post operatively.

- Focus on high quality proteins such as poultry, lean beef, eggs, nuts, seeds, fish.
- Consider an oral nutrition supplement/drink for additional protein intake.



# Pre-Surgery: Optimization

Research has shown that taking an immunonutrition supplement before and after surgery improves healing (take as directed on container).

There are several immunonutrition supplements available, including:

- Nestle Impact Advanced Recovery (Impact AR) can be purchased on Amazon, Walmart.com, or at the Nestle Store at 1-888-240-2713
- Ensure Surgery can be purchased on Amazon or Walmart.com
- Another option is to ask your local pharmacy if they can order it for you

**If you are a diabetic**, please talk to the physician managing your diabetes before using or purchasing Impact AR or Ensure Surgery.



For more information go to: <https://www.facs.org/Quality-Programs/strong-for-surgery>



# Pre-Surgery: Optimization

<b>Stop Smoking</b>	Stop smoking now. <ul style="list-style-type: none"><li>• 4 Weeks before surgery reduces complications 20-30%.</li></ul>
<b>Stop Alcohol</b>	Stop alcohol 1 week before surgery.
<b>No Dental Work</b>	No dental work 2 weeks before surgery. <ul style="list-style-type: none"><li>• Discuss with your surgeon and dentist the guidelines for antibiotics with all future dental care.</li></ul>
<b>No Manicures or Pedicures</b>	No professional manicure/pedicure 2 weeks before surgery.
<b>No Shaving</b>	No shaving any body hair below the neck the 3 days prior to your surgery. <ul style="list-style-type: none"><li>• Facial shaving is permitted before surgery.</li></ul>



# Pre-Surgery: Medications to Stop

Consult your prescribing physician for when to **stop** a PRESCRIBED blood thinner. **NSAIDs** need to be **stopped 7 days** prior to surgery.

## Prescription Blood Thinner Examples

- Aspirin
- Coumadin
- Eliquis
- Lixiana
- Plavix
- Pradaxa
- Xarelto



## NSAID Examples

- Aspirin (Bufferin, Ecotrin, Aspercreme...)
- Aspirin containing drugs
- Ibuprofen – Advil, Motrin, Nuprin
- Naproxen – Aleve
- Voltaren – diclofenac
- Mobic – meloxicam
- Celebrex – celecoxib
- Indomethacin

# Pre-Surgery: GLP-1 agonist Medications

If you are taking a GLP-1 agonist medication for weight loss, talk to your surgeon. These medications often need to be stopped before surgery.

## Examples:

- Dulaglutide (Trulicity)
- Exenatide extended release (Bydureon bcise)
- Exenatide (Byetta)
- Semaglutide (Ozempic)
- Liraglutide (Victoza, Saxenda)



# Pre-Surgery: Stopping Supplements

## Stop taking herbal supplements 14 days before surgery

Herbal supplements are derived from different parts of plant.

They come in many forms including:

- Tablets, capsules, teas, tinctures, oils, lotions, or ointments

These are a few examples:

- CBD or THC containing products, aloe vera, echinacea, ephedra, feverfew, green tea, garlic, ginkgo biloba, ginseng, ginger, golden seal, kava, licorice, saw palmetto, Valeria root, Saint John's wort, turmeric

Discuss any other supplements or vitamins that you take with your surgeon.

- Your surgeon will notify you if they need to be stopped.



## Infection Prevention: Before and After Surgery

Wash your hands often with soap and water or an alcohol-based hand sanitizer.

Avoid touching your eyes, nose and mouth; germs spread this way.

Brush your teeth.


Avoid contact with sick people.

Clean and disinfect surfaces and objects that may be contaminated with germs.



# Pre-Surgery: Infection Prevention

Your surgeon will give you directions for showering with **Chlorhexidine gluconate (CHG)**



**CHG** is both a soap and antiseptic that kills germs on contact.

- **CHG** safely binds to the skin and can continue killing germs for up to 24 hours.
- **CHG** has a cumulative effect, so the protection against germs increases with repeated use.

## **Nasal antiseptic**

- **Nasal antiseptic** has been shown to inhibit the growth of bacteria in the nose over a period of up to 12 hours in studies.

**Day of Surgery:** Do not apply lotion, creams, powder, or makeup on the day of surgery.

# Pre-Surgery: The Day Before Surgery

Review instructions provided by the surgeon's office.

Make sure you know the time to stop eating and drinking.

Know the time to be at the surgery center and please be on time.

Infection Prevention, the night before surgery

- Wear clean pajamas
- Sleep on clean sheets
- Keep pets off your bed



# Pre-Surgery: Hydration

## THE NIGHT BEFORE SURGERY

Drink one of these the night before your surgery IF recommended by your surgeon:

- 16fl oz (2cups) Gatorade

**OR**

- 2 Bottles Ensure® Pre-Surgery Carbohydrate **Clear** Nutrition Drink

**DO NOT eat any solid food after midnight**

## THE DAY OF SURGERY


- You may drink CLEAR liquids up to 3 hours before your scheduled surgery time, IF okayed by your surgeon.
- Limit the clear liquid volume to 1½ cups or 12 ounces per hour.

Drink one of these 2-3 hours before your surgery, IF recommended by your surgeon:

- 16fl oz (2cups) Gatorade

**OR**

- 1 Bottle Ensure® Pre-Surgery Carbohydrate **Clear** Nutrition Drink



# Pre-Surgery: Hydration

What clear liquids can I drink and NOT drink the morning of surgery?

Allowed	DO NOT CONSUME
Water	Milk or Dairy Products
Apple & Cranberry Juice	Citrus Juices
Gatorade or equivalent carb containing sports drinks	Prune Juice
Ensure® Pre-Surgery Carbohydrate <u>Clear</u> Nutrition Drink	Juices with Pulp
<u>Plain</u> Coffee or Tea (No milk or creamer)	Alcoholic Beverages

Individuals with obesity, diabetes, gastroparesis, or any known delayed stomach emptying issues are not to drink liquids after midnight.

Follow instructions carefully or your surgery may be canceled.



# Day of Surgery: Pre-Op

Consents reviewed and signed.

An intravenous line will be placed.

You will meet your Anesthesiologist and discuss anesthesia.

- Most patients have spinal anesthesia
  - Patients are asleep with spinal anesthesia



# Day of Surgery: Operating Room

Anesthetic medication will be given.

You will be positioned on a table that is specially designed for your surgery.

Your surgical area will be scrubbed with surgical prep.



# Post Anesthesia Care Unit (PACU)

You will “wake up” in the PACU (recovery room)

Your nurse:

- Assesses your medical condition
- Provides pain management as needed
- Teaches you to get in and out of bed
- Instructs you on how to walk with a walker

Before discharge:


- X-rays are taken of your new joint
- You will urinate



# Post Op Precautions

## Posterior Total Hip Precautions

- No bending past 90 degrees
- No crossing your legs
- No internal rotation



\*These precautions are in place until your surgeon clears you from them.

# After Discharge: Physical Therapy

## Physical therapy after surgery

- Arrangements are made by the surgeon office for **home health** physical therapy, if your surgeon determines it is needed.
- Your surgeon will give you a prescription for **outpatient** physical therapy when you have been cleared to start it.



# At Home: Preventing Pain

## Cold Therapy

Use ice packs or a cold therapy ice machine.  
Never apply ice therapy directly to the skin. Have a protective barrier such as a towel.  
Check your skin often.  
Allow time for your skin to warm to its normal temperature between applications.

## Elevation

Think “toes above the nose.”  
Elevate the affected extremity above the level of the heart if possible.  
Elevation reduces swelling which in turn reduces pain.

## Distraction

Distract yourself with relaxing activities you can do while sitting or laying down.

- TV, electronics, reading

## Oral Medications

Take only as prescribed.  
PRN means as needed. Do not take PRN medication if you do not need it.  
Narcotics -use the minimal dose of the narcotic to get the maximum effect.

# At Home: Activity, the First Few Weeks

## Lying Down & Elevating the Surgical Leg

“Toes above the nose.”

Elevate 3-4 times a day for 45 minutes to an hour during the day.  
Remember your precautions.



## Sitting

Keep your knees below the level of the hip. Remember precautions.  
Sit leaning slightly back with your feet elevated on a stool or equivalent to help prevent swelling. (Think first class on an airplane)  
Limit sitting with your feet on the ground to meals and chair exercises to avoid getting stiff.



## Walking

Take short walks in the house hourly while awake.  
Use your walker or cane as determined by your therapist.




- Remember to move / wiggle around when lying down and sitting. Pump your feet often. This will help to keep you comfortable.

# At Home: Deep Breathing Exercises

## Deep Breathing

1. Sit up straight and tall
2. Relax
3. Take a deep breath, slow breath in
4. Hold your breath in for 3 seconds
5. Let it out
6. Repeat about 10 breaths every hour, but not 10 times in a row





# At Home: DVT Prevention


A DVT (deep vein thrombosis) is a rare surgical complication. It is important to know the signs.

## Let your surgeon know if you:

- Have new swelling in one leg not related to an injury
- Your calf (back of lower leg) is tender or painful when you push on it
- Your calf feels warm or hot to touch compared to the other leg

## Prevent Blood Clots:

- Get moving
  - Take short, frequent walks
  - Perform ankle pumps when sitting or lying down
    - at least 10 times an hour



# At Home: DVT Prevention, Blood Thinner

You will be placed on a blood thinner after surgery.

Know your blood thinner.

Your blood thinner must be taken as prescribed.

Aspirin is a commonly prescribed as a blood thinner after surgery.



# At Home: Recovery

## Expect:

### Pain

- Take it easy the first two weeks
- Use your pain medications as prescribed

### Swelling

- Elevate
- Ice

### Bruising

- Bruising is normal
- Bruising can go down the whole leg after total hip surgery

## Be Aware:

- Dizziness / Lightheadedness can occur
- Go slow when standing up
- Use your walker

# At Home: When to call the Surgeon

## Call your surgeon if you have any:

- Severe pain that is not controlled with pain medicine or is getting worse
- Heavy bleeding from the incision
- Signs of infection:
  - Fever over 102 degrees Fahrenheit
  - Redness, swelling, warmth around the incision or drainage of pus
  - Severe vomiting or cannot keep food down

# Thank You



## Getting You Back to You

# Hoag Orthopedic Institute

The information in this presentation is designed as a general guide, while the information provided by your physician is specific to your individual needs.

